


A




MTANNDDDIK SPIPITNK TL SQL **KRFERS** GRPSSSQGEI **KRKKS**R LYAA DGRPHSPLR A
RASTPTLQDQ KLFNGMDSTS LLNERLQHYT LDYVSDR AQH MKNIDPSSR WFSRSVRPEF
 PIEEFLPYKT ESHEDQAKYL CHVLVNLVIA ISSLDIQGLI SISSKDLADL KKEVDDLALK
 TDLFRLSNNT AENDLLGNDI ADYDDAEGLE DELDEYFDLA GPDFNATGKI TAKSATIVNV
 NHWTNELKNC LHFDFPVALR KSLATVYYYL SLVQGQKVYR QMHVDMFER L VSLDDDRTNF
TELLQKQGLL LDHQIMLNFL CEFLPYDPDP YARYELSSKE DLQLFRLLK HAHNAKPFFD
 KSKESLLVDT MNFLSSLAP STMMAVMPIV TSVVPYHYHI HSKIIDYFPF CYSIWSSVSA
 NVAIDTHMYD FVGSISKDVH NKILSSEHEK DVVGVEFGEF GIFTDDQMTF MFNRLQGHLR
 TDGQIHSYSR TVKPFVYAIN GSKKDRFEK LVSLAKAIET FIHPSNNGFW TKPNAKFVHA
 FIKSYHGRVK YEEDICARGV TNGICLTSFC HEEIVEIFLN IISLGSQKNK PDIANYYISC
 FAYLLELDPS NAYLIYDKIL IDLYDTLADQ FINSRHRIIS SLKQFTRVIR FIVMDKLYRV
 HITNVLSMLV SKLDMNDTNL TSNLINGIVS IAAFIPIQDL TGEDDYISFE SDTLPVQQH
 FYHIKCGESS KTRFVDELL NNAFKASTTV FQSMKVYVE KIFQLVDVDL EDSLVTKINQ
 TTMILQESMD DKIFYNFASL LQRNFSNDS FKEKDPNYEL VTIPLAALVR RNNGLSKELV
 RTLLFHIKEQ IKGAGSVRS TSEIQQRDVK LVLYLTALND VLRQCHESLL EYSEDLITFM
 KYLYDNVTNP PLDVITSIVI HSALATLCTT EITDCRLFPE DSKIPEKDR W GGLQFDPR RF
 DKQHLSFQWH VPSSDEITLS ISILESLEY CINNVEELMK APRHDSEYGD MIQKYVLVMT
 HTLSGSSLLF DPDFNKYRTQ SNLSYREKLI LLKNIRENNC DPQELDIDIE QIR SGKDD
YIESKDIEAG LNAGVSDVVQ LRDEFDEL I VDEEVVSEMP SGVNTPIAGT HGTDNSAMSS
 DLAFRDLDIY TCNYYFGNTT EEKLQNPQYL QVHRVRARIG HFFHKLYVFL STNFENNTNM
 FQILLHGLKV WFTDLGQETV FNEDPNAFID VDFLENVQSL SHVNEPFTRT NFAIRANSLH
 QSRVLLHSTN RKASKLENLL LVDIIQLATS LYPDIYKPAQ GTLVHCKMQL VGSYGVVINK
 IIPSLEKAIK DHDYMKIQVI LNVLLIKKIH RKLMTDYKDI GRLIFLLIEC CRVNELEIGM
 YADKILTDIV IGIKIPSSVC VISDQAFPL APPDGTINLQ VEAVKLAKKK KREYYLSLLV
 DLQDKLLDKL DNEKDMGWKI RMFILRFVTQ IQSNLESKPD KRAVFSIISQ ISTKHPEIHH
 LVVKSLLSTC NKIISLSDYE YDITRAYKNE FNPSFVEILD TSTTSFPKTF TEEMNFDNP
 KYFIDLRAYV GWLCWGRMY VMSPKALKLN LR ENELEVLK TAGHLLTREF LR DVTMNLVQ
DNETRGVFSS GNVSFFSLVI LLISSGFCEL NMSDLFELCE SYYNKDDKAS MIMSVEIVAG
 LVCGSKFMSV SLDLKRDTFI ENFLAKCLDY ELNHDAFEIW STLAWWLPV VDLRRSKTFF
 CHFINADGMF DRESAATHQ TSKIYMLRSI LMSMEFRAPD V GK LFDELVF DHPYDQVR QA
 VAKLLTTLVQ NQSNPSISDP TTLLEAERND PDGLGLPLKS VPEKVDAYIK KQFEIKNLE
 DSVVGLNPQQ FIKTDYFYRT STIFYWIKEM ARGPNKVLLV PYLVDYVLPF LIGLVKHKDV
 CALASLDPVR LYAGLGYPYI RKNHVAAIVD YVCSSNVALS SNQTKLQLAF IQHFLSAELL
 QLTEEEKNKI LEFVVSNNLYN EQFVEVRVRA ASILSDIVHN WKEEQPLLSL IERFAKGLDV
 NKYTSKERQK LSKTDIKIHG NVLGLGAIIS AFPYVFPLPP WIPK QLSNLS SWAR TSGMTG
 QAAKNTISEF KKVRA DTWK F DRASFNT EEL EDLEGVLWRS YYA

Figure S1 (A) Amino acid sequence of Blm3 as deduced from the BLM3-YFL006 gene of yeast *Saccharomyces cerevisiae*. The putative intergenic region between BLM3 and YFL006 was sequenced. BLM3 was confirmed to be in frame with YFL006 (Yeast Proteome Database). The N-terminal peptide obtained by Edman degradation and tryptic peptides from Blm3 are underlined. The putative nuclear localization sequence is highlighted in grey. The protein kinase phosphorylation site is framed (Ficarro et al. (2002). Phosphoproteome analysis by mass spectrometry and its application to *Saccharomyces cerevisiae*. Nat. Biotechnol.




Ms 81206
Figure S1B

(B) The *blm3D* null strain is temperature sensitive. Growth curves of *blm3D* compared to isogenic wild type cells in YPD at 28°C and 37°C. Spot assays of *blm3D* compared to isogenic wild type cells grown on YPD plates at 28°C and 37°C.




Ms 81206
Figure S1C

(C) Galactose-induced overexpression of Blm3 inhibits cell growth. To express Blm3 behind the GAL1 promoter, the endogenous BLM3 promoter (-60bp referred to ATG) was chromosomally disrupted by the GAL1 promoter. This was achieved by chromosomal integration of LEU2-GAL1 flanked by homologous sequences of the BLM3 promoter region. On galactose-containing media wild type cells expressing Blm3 behind the GAL1 promoter were growth-inhibited compared to those expressing Blm3 behind the endogenous promoter (upper panels). Overexpression of Blm3 was verified by Western blot analysis using anti-Blm3 antibodies (lower panel; wild type cells expressing Blm3 behind the endogenous promoter (lane 1) and behind the GAL1 promoter (lane 2)). Overexpression of Blm3 did not cause an accumulation of precursor complexes (not shown) indicating that the n-CP rather than Blm3 are limiting in late maturation processes.



Ms 81206
Figure S2

Figure S2. Blm3 co-migrates with Ump1 in density gradient ultracentrifugation. (A) Total extracts of wild type cells expressing Ump1-HA instead of the endogenous protein were subjected to 10-40% glycerol gradient ultracentrifugation as previously described (Enenkel et al., 1998). Fractions were collected from top to bottom and assayed for peptide cleavage activity against Cbz-Leu-Leu-Glu-b-naphthylamide (upper panel). Protein samples of each fraction were run on SDS-PAGE, blotted and probed for Blm3 and Ump1-HA. (B) Extracts of wild type cells expressing GFP-tagged Blm3 were subjected to 10 - 40% glycerol gradient ultracentrifugation as described above and probed for GFP-tagged Blm3.



Ms 81206
Figure S3

Figure S3. (A) GFP-labelled Blm3 is nuclear in *ump1D* cells expressing Blm3-GFP instead of the endogenous protein. Cells were monitored by direct fluorescence microscopy using a GFP filter (left panel) and Nomarski optics (right panel). Precursor complexes were found to be predominantly nuclear in *ump1D* cells and to be associated with Blm3 (not shown). (B) GFP-labelled Ump1 and Pre6 (a4) are mainly nuclear in *blm3D* cells. *blm3D* (left panels) and isogenic wild type cells (right panels) expressing either GFP-tagged Ump1 or Pre6 (a4) were grown to logarithmic phase at 28°C and shifted to 37°C for 4 hours (left panel, GFP filter; right panel, Nomarski optics). Homologous recombination techniques were used to replace proteasomal proteins by GFP-tagged versions in *blm3D* and *ump1D* cells (Enenkel et al., 1998).





Table I: Inhibitory effect of Blm3 on m-CP activity

Substrate	Proteasomal activity [%±10]
Z-Gly-Gly-Leu-AMC	114
Succ-Leu-Leu-Val-Tyr-AMC	36
Z-Leu-Leu- Glu-βNA	79
<hr/>	
Lactacystin (LLVY))	0
Regulatory 19S particle (LLVY)	250

B In vivo degradation of FBPase



Ms
Fig

Figure S4. Inhibitory effect of Blm3 on m-CP activity. (A) SDS-PAGE followed by Coomassie blue staining of purified Blm3. Blm3 was dissociated from affinity-purified proteasomal complexes by 600 mM NaCl and transferred into assay buffer. Table I: Proteasomal activity was measured in 20 mM Tris/HCl pH 7.8, 5 mM MgCl₂, 10 mM KCl, 1 mM DTT for 30 min at 37°C using 100 μM chromogenic substrates Z-LLE-b-naphthylamide (NA), Z-GGL-7-amido-4-methylcoumarin (AMC) and Succ-LLVY-AMC (Ustrell et al., 2002). All measurements were referred to the value of the m-CP without Blm3 (100 %). Complete inhibition of the m-CP was achieved by 50 μM lactacystin. CP and Blm3 were incubated in a molar ratio of 1:10 (70 fmol m-CP). RP was purified (Saeki, Y., Toh-e, A. and Yokosawa, H. (2000) Rapid Isolation and Characterization of the Yeast Proteasome Regulatory Complex. Biochem. Biophys. Res. Commun., 273, 509-515) and added in equimolar ratio. Purified Blm3 had no proteasomal peptide cleavage activity. (B) In vivo degradation of the proteasomal substrate fructose-bisphosphatase (FBPase) in wild type and *blm3D* cells as analyzed by cycloheximide chase experiments (Horak, J., Regelmann, J., Wolf, D. H. (2002) Two distinct proteolytic systems responsible for glucose-induced degradation of fructose-1,6-bisphosphatase and the Gal2p transporter in the yeast *Saccharomyces cerevisiae* share the same protein components of the glucose signaling pathway. J Biol Chem 277(10): 8248-8254). We thank the lab members of Dieter Wolf (University of Stuttgart) for support in this experiment.